

Bayesian Structural Equation Modeling (Methodology in the Social Sciences)

Get your blood sugar tested if you have any of the symptoms of diabetes.

If you have any of the following diabetes symptoms, see your doctor about getting your blood sugar tested:

Urinate (pee) a lot, often at night

Are very thirsty

Lose weight without trying

Are very hungry

Have blurry vision

Have numb or tingling hands or feet

Feel very tired

Have very dry skin

Have sores that heal slowly

Have more infections than usual

Symptoms of Type 1 Diabetes

People who have type 1 diabetes may also have nausea, vomiting, or stomach pains. Type 1 diabetes symptoms can develop in just a few weeks or months and can be severe. Type 1 diabetes usually starts when you're a child, teen, or young adult but can happen at any age.

Symptoms of Type 2 Diabetes

Type 2 diabetes symptoms often take several years to develop. Some people don't notice any symptoms at all. Type 2 diabetes usually starts when you're an adult, though more and more children and teens are developing it. Because symptoms are hard to spot, it's important to know the risk factors for type 2 diabetes. Make sure to visit your doctor if you have any of them.

Symptoms of Gestational Diabetes

Gestational diabetes (diabetes during pregnancy) usually doesn't have any symptoms. If you're pregnant, your doctor should test you for gestational diabetes between 24 and 28 weeks of pregnancy. If needed, you can make changes to protect your health and your baby's health.

Reference

[Nursing: A Concept-Based Approach to Learning, Volume 2](#)

[Exploring Protein Structure: Principles and Practice \(Learning Materials in Biosciences\)](#)