

Case Files Obstetrics and Gynecology, Fifth Edition

Recommendations

The recommendations are grouped into three areas for consideration:

Determining when to initiate or continue opioids for chronic pain.

Opioid selection, dosage, duration, follow-up, and discontinuation.

Assessing risk and addressing harms of opioid use.

There are 12 recommendations (Box 1). Each recommendation is followed by a rationale for the recommendation, with considerations for implementation noted. In accordance with the ACIP GRADE process, CDC based the recommendations on consideration of the clinical evidence, contextual evidence (including benefits and harms, values and preferences, resource allocation), and expert opinion. For each recommendation statement, CDC notes the recommendation category (A or B) and the type of the evidence (1, 2, 3, or 4) supporting the statement (Box 2). Expert opinion is reflected within each of the recommendation rationales. While there was not an attempt to reach consensus among experts, experts from the Core Expert Group and from the Opioid Guideline Workgroup (“experts”) expressed overall, general support for all recommendations. Where differences in expert opinion emerged for detailed actions within the clinical recommendations or for implementation considerations, CDC notes the differences of opinion in the supporting rationale statements.

Category A recommendations indicate that most patients should receive the recommended course of action; category B recommendations indicate that different choices will be appropriate for different patients, requiring clinicians to help patients arrive at a decision consistent with patient values and preferences and specific clinical situations. Consistent with the ACIP (47) and GRADE process (48), category A recommendations were made, even with type 3 and 4 evidence, when there was broad agreement that the advantages of a clinical action greatly outweighed the disadvantages based on a consideration of benefits and harms, values and preferences, and resource allocation. Category B recommendations were made when there was broad agreement that the advantages and disadvantages of a clinical action were more balanced, but advantages were significant enough to warrant a recommendation. All recommendations are category A recommendations, with the exception of recommendation 10, which is rated as category B. Recommendations were associated with a range of evidence types, from type 2 to type 4.

In summary, the categorization of recommendations was based on the following assessment:

No evidence shows a long-term benefit of opioids in pain and function versus no opioids for chronic pain with outcomes examined at least 1 year later (with most placebo-controlled randomized trials ≥ 6 weeks in duration).

Extensive evidence shows the possible harms of opioids (including opioid use

P

disorder, overdose, and motor vehicle injury).

Extensive evidence suggests some benefits of nonpharmacologic and nonopioid pharmacologic treatments compared with long-term opioid therapy, with less harm.

Determining When to Initiate or Continue Opioids for Chronic Pain

1. Nonpharmacologic therapy and nonopioid pharmacologic therapy are preferred for chronic pain. Clinicians should consider opioid therapy only if expected benefits for both pain and function are anticipated to outweigh risks to the patient. If opioids are used, they should be combined with nonpharmacologic therapy and nonopioid pharmacologic therapy, as appropriate (recommendation category: A, evidence type: 3).

Patients with pain should receive treatment that provides the greatest benefits relative to risks. The contextual evidence review found that many nonpharmacologic therapies, including physical therapy, weight loss for knee osteoarthritis, psychological therapies such as CBT, and certain interventional procedures can ameliorate chronic pain. There is high-quality evidence that exercise therapy (a prominent modality in physical therapy) for hip (100) or knee (99) osteoarthritis reduces pain and improves function immediately after treatment and that the improvements are sustained for at least 24 months. Previous guidelines have strongly recommended aerobic, aquatic, and/or resistance exercises for patients with osteoarthritis of the knee or hip (176). Exercise therapy also can help reduce pain and improve function in low back pain and can improve global well-being and physical function in fibromyalgia (98,101). Multimodal therapies and multidisciplinary biopsychosocial rehabilitation-combining approaches (e.g., psychological therapies with exercise) can reduce long-term pain and disability compared with usual care and compared with physical treatments (e.g., exercise) alone. Multimodal therapies are not always available or reimbursed by insurance and can be time-consuming and costly for patients. Interventional approaches such as arthrocentesis and intraarticular glucocorticoid injection for pain associated with rheumatoid arthritis (117) or osteoarthritis (118) and subacromial corticosteroid injection for rotator cuff disease (119) can provide short-term improvement in pain and function. Evidence is insufficient to determine the extent to which repeated glucocorticoid injection increases potential risks such as articular cartilage changes (in osteoarthritis) and sepsis (118). Serious adverse events are rare but have been reported with epidural injection (120).

Several nonopioid pharmacologic therapies (including acetaminophen, NSAIDs, and selected antidepressants and

anticonvulsants) are effective for chronic pain. In particular, acetaminophen and NSAIDs can be useful for arthritis and low back pain. Selected anticonvulsants such as pregabalin and gabapentin can improve pain in diabetic neuropathy and post-herpetic neuralgia (contextual evidence review). Pregabalin, gabapentin, and carbamazepine are FDA-approved for treatment of certain neuropathic pain conditions, and pregabalin is FDA approved for fibromyalgia management. In patients with or without depression, tricyclic antidepressants and SNRIs provide effective analgesia for neuropathic pain conditions including diabetic neuropathy and post-herpetic neuralgia, often at lower dosages and with a shorter time to onset of effect than for treatment of depression (see contextual evidence review). Tricyclics and SNRIs can also relieve fibromyalgia symptoms. The SNRI duloxetine is FDA-approved for the treatment of diabetic neuropathy and fibromyalgia. Because patients with chronic pain often suffer from concurrent depression (144), and depression can exacerbate physical symptoms including pain (177), patients with co-occurring pain and depression are especially likely to benefit from antidepressant medication (see Recommendation 8). Nonopioid pharmacologic therapies are not generally associated with substance use disorder, and the numbers of fatal overdoses associated with nonopioid medications are a fraction of those associated with opioid medications (contextual evidence review). For example, acetaminophen, NSAIDs, and opioid pain medication were involved in 881, 228, and 16,651 pharmaceutical overdose deaths in the United States in 2010 (178). However, nonopioid pharmacologic therapies are associated with certain risks, particularly in older patients, pregnant patients, and patients with certain co-morbidities such as cardiovascular, renal, gastrointestinal, and liver disease (see contextual evidence review). For example, acetaminophen can be hepatotoxic at dosages of > 3-4 grams/day and at lower dosages in patients with chronic alcohol use or liver disease (109). NSAID use has been associated with gastritis, peptic ulcer disease, cardiovascular events (111,112), and fluid retention, and most NSAIDs (choline magnesium trisilicate and selective COX-2 inhibitors are exceptions) interfere with platelet aggregation (179). Clinicians should review FDA-approved labeling including boxed warnings before initiating treatment with any pharmacologic therapy.

Although opioids can reduce pain during short-term use, the clinical evidence review found insufficient evidence to determine whether pain relief is sustained and whether function or quality of life improves with long-term opioid therapy (KQ1). While benefits for pain relief, function, and quality of life with long-term opioid use for chronic pain are uncertain, risks associated with long-term opioid use are clearer and significant. Based on the clinical evidence review, long-term opioid use for chronic pain is associated with serious risks including increased risk for opioid use disorder, overdose, myocardial infarction, and motor vehicle injury (KQ2). At a population level, more than 165,000 persons in the United States have died from opioid pain-medication-related overdoses since 1999 (see Contextual Evidence Review).

Integrated pain management requires coordination of medical, psychological, and social aspects of health care and includes primary care, mental health care, and specialist services when needed (180). Nonpharmacologic physical and psychological treatments such as exercise and CBT are approaches that encourage active patient participation in the care plan, address the effects of pain in the patient's life, and can result in sustained improvements in pain and function without apparent risks. Despite this, these therapies are not always or fully covered by insurance, and access and cost can be barriers for patients. For many patients, aspects of these approaches can be used even when there is limited access to specialty care. For example, previous guidelines have strongly recommended aerobic, aquatic, and/or resistance exercises for patients with osteoarthritis of the knee or hip (176) and maintenance of activity for patients with low back pain (110). A randomized trial found no difference in reduced chronic low back pain intensity, frequency or disability between patients assigned to relatively low-cost group aerobics and individual physiotherapy or muscle reconditioning sessions (181). Low-cost options to integrate exercise include brisk walking in public spaces or use of public recreation facilities for group exercise. CBT addresses psychosocial contributors to pain and improves function (97). Primary care clinicians can integrate elements of a cognitive behavioral approach into their practice by encouraging patients to take an active role in the care plan, by supporting patients in engaging in beneficial but potentially anxiety-provoking activities, such as exercise (179), or by providing education in relaxation techniques and coping strategies. In many locations, there are free or low-cost patient support, self-help, and educational community-based programs that can provide stress reduction and other mental health benefits. Patients with more entrenched anxiety or fear related to pain, or other significant psychological distress, can be referred for formal therapy with a mental health specialist (e.g., psychologist, psychiatrist, clinical social worker). Multimodal therapies should be considered for patients not responding to single-modality therapy, and combinations should be tailored depending on patient needs, cost, and convenience.

To guide patient-specific selection of therapy, clinicians should evaluate patients and establish or confirm the diagnosis. Detailed recommendations on diagnosis are provided in other guidelines (110,179), but evaluation should generally include a focused history, including history and characteristics of pain and potentially contributing factors (e.g., function, psychosocial stressors, sleep) and physical exam, with imaging or other diagnostic testing only if indicated (e.g., if severe or progressive neurologic deficits are present or if serious underlying conditions are suspected) (110,179). For complex pain syndromes, pain specialty consultation can be considered to assist with diagnosis as well as management. Diagnosis can help identify disease-specific interventions to reverse or ameliorate pain; for example, improving glucose control to prevent progression of diabetic neuropathy; immune-modulating agents for rheumatoid arthritis; physical or occupational therapy to address posture, muscle weakness, or repetitive

occupational motions that contribute to musculoskeletal pain; or surgical intervention to relieve mechanical/compressive pain (179). The underlying mechanism for most pain syndromes can be categorized as neuropathic (e.g., diabetic neuropathy, postherpetic neuralgia, fibromyalgia), or nociceptive (e.g., osteoarthritis, muscular back pain). The diagnosis and pathophysiologic mechanism of pain have implications for symptomatic pain treatment with medication. For example, evidence is limited or insufficient for improved pain or function with long-term use of opioids for several chronic pain conditions for which opioids are commonly prescribed, such as low back pain (182), headache (183), and fibromyalgia (184). Although NSAIDs can be used for exacerbations of nociceptive pain, other medications (e.g., tricyclics, selected anticonvulsants, or transdermal lidocaine) generally are recommended for neuropathic pain. In addition, improvement of neuropathic pain can begin weeks or longer after symptomatic treatment is initiated (179). Medications should be used only after assessment and determination that expected benefits outweigh risks given patient-specific factors. For example, clinicians should consider falls risk when selecting and dosing potentially sedating medications such as tricyclics, anticonvulsants, or opioids, and should weigh risks and benefits of use, dose, and duration of NSAIDs when treating older adults as well as patients with hypertension, renal insufficiency, or heart failure, or those with risk for peptic ulcer disease or cardiovascular disease. Some guidelines recommend topical NSAIDs for localized osteoarthritis (e.g., knee osteoarthritis) over oral NSAIDs in patients aged ≥ 75 years to minimize systemic effects (176).

Experts agreed that opioids should not be considered first-line or routine therapy for chronic pain (i.e., pain continuing or expected to continue >3 months or past the time of normal tissue healing) outside of active cancer, palliative, and end-of-life care, given small to moderate short-term benefits, uncertain long-term benefits, and potential for serious harms; although evidence on long-term benefits of nonopioid therapies is also limited, these therapies are also associated with short-term benefits, and risks are much lower. This does not mean that patients should be required to sequentially “fail” nonpharmacologic and nonopioid pharmacologic therapy before proceeding to opioid therapy. Rather, expected benefits specific to the clinical context should be weighed against risks before initiating therapy. In some clinical contexts (e.g., headache or fibromyalgia), expected benefits of initiating opioids are unlikely to outweigh risks regardless of previous nonpharmacologic and nonopioid pharmacologic therapies used. In other situations (e.g., serious illness in a patient with poor prognosis for return to previous level of function, contraindications to other therapies, and clinician and patient agreement that the overriding goal is patient comfort), opioids might be appropriate regardless of previous therapies used. In addition, when opioid pain medication is used, it is more likely to be effective if integrated with nonpharmacologic therapy. Nonpharmacologic approaches such as exercise and CBT should be used to reduce pain and improve function in patients with chronic pain.

Nonopioid pharmacologic therapy should be used when benefits outweigh risks and should be combined with nonpharmacologic therapy to reduce pain and improve function. If opioids are used, they should be combined with nonpharmacologic therapy and nonopioid pharmacologic therapy, as appropriate, to provide greater benefits to patients in improving pain and function.

2. Before starting opioid therapy for chronic pain, clinicians should establish treatment goals with all patients, including realistic goals for pain and function, and should consider how opioid therapy will be discontinued if benefits do not outweigh risks. Clinicians should continue opioid therapy only if there is clinically meaningful improvement in pain and function that outweighs risks to patient safety (recommendation category: A, evidence type: 4).

The clinical evidence review found insufficient evidence to determine long-term benefits of opioid therapy for chronic pain and found an increased risk for serious harms related to long-term opioid therapy that appears to be dose-dependent. In addition, studies on currently available risk assessment instruments were sparse and showed inconsistent results (KQ4). The clinical evidence review for the current guideline considered studies with outcomes examined at 1 year that compared opioid use versus nonuse or placebo. Studies of opioid therapy for chronic pain that did not have a nonopioid control group have found that although many patients discontinue opioid therapy for chronic noncancer pain due to adverse effects or insufficient pain relief, there is weak evidence that patients who are able to continue opioid therapy for at least 6 months can experience clinically significant pain relief and insufficient evidence that function or quality of life improves (185). These findings suggest that it is very difficult for clinicians to predict whether benefits of opioids for chronic pain will outweigh risks of ongoing treatment for individual patients. Opioid therapy should not be initiated without consideration of an "exit strategy" to be used if the therapy is unsuccessful.

Experts agreed that before opioid therapy is initiated for chronic pain outside of active cancer, palliative, and end-of-life care, clinicians should determine how effectiveness will be evaluated and should establish treatment goals with patients. Because the line between acute pain and initial chronic pain is not always clear, it might be difficult for clinicians to determine when they are initiating opioids for chronic pain rather than treating acute pain. Pain lasting longer than 3 months or past the time of normal tissue healing (which could be substantially shorter than 3 months, depending on the condition) is generally no longer considered acute. However, establishing treatment goals with a patient who has already received opioid therapy for 3 months would defer this discussion well past the point of

initiation of opioid therapy for chronic pain. Clinicians often write prescriptions for long-term use in 30-day increments, and opioid prescriptions written for 30 days are likely to represent initiation or continuation of long-term opioid therapy. Before writing an opioid prescription for 30 days, clinicians should establish treatment goals with patients. Clinicians seeing new patients already receiving opioids should establish treatment goals for continued opioid therapy. Although the clinical evidence review did not find studies evaluating the effectiveness of written agreements or treatment plans (KQ4), clinicians and patients who set a plan in advance will clarify expectations regarding how opioids will be prescribed and monitored, as well as situations in which opioids will be discontinued or doses tapered (e.g., if treatment goals are not met, opioids are no longer needed, or adverse events put the patient at risk) to improve patient safety.

Experts thought that goals should include improvement in both pain relief and function (and therefore in quality of life). However, there are some clinical circumstances under which reductions in pain without improvement in physical function might be a more realistic goal (e.g., diseases typically associated with progressive functional impairment or catastrophic injuries such as spinal cord trauma). Experts noted that function can include emotional and social as well as physical dimensions. In addition, experts emphasized that mood has important interactions with pain and function. Experts agreed that clinicians may use validated instruments such as the three-item Pain average, interference with Enjoyment of life, and interference with General activity (PEG) Assessment Scale (186) to track patient outcomes. Clinically meaningful improvement has been defined as a 30% improvement in scores for both pain and function (187). Monitoring progress toward patient-centered functional goals (e.g., walking the dog or walking around the block, returning to part-time work, attending family sports or recreational activities) can also contribute to the assessment of functional improvement. Clinicians should use these goals in assessing benefits of opioid therapy for individual patients and in weighing benefits against risks of continued opioid therapy (see Recommendation 7, including recommended intervals for follow-up). Because depression, anxiety, and other psychological co-morbidities often coexist with and can interfere with resolution of pain, clinicians should use validated instruments to assess for these conditions (see Recommendation 8) and ensure that treatment for these conditions is optimized. If patients receiving opioid therapy for chronic pain do not experience meaningful improvements in both pain and function compared with prior to initiation of opioid therapy, clinicians should consider working with patients to taper and discontinue opioids (see Recommendation 7) and should use nonpharmacologic and nonopioid pharmacologic approaches to pain management (see Recommendation 1).

3. Before starting and periodically during opioid therapy, clinicians should discuss with patients known risks and

realistic benefits of opioid therapy and patient and clinician responsibilities for managing therapy (recommendation category: A, evidence type: 3).

The clinical evidence review did not find studies evaluating effectiveness of patient education or opioid treatment plans as risk-mitigation strategies (KQ4). However, the contextual evidence review found that many patients lack information about opioids and identified concerns that some clinicians miss opportunities to effectively communicate about safety. Given the substantial evidence gaps on opioids, uncertain benefits of long-term use, and potential for serious harms, patient education and discussion before starting opioid therapy are critical so that patient preferences and values can be understood and used to inform clinical decisions. Experts agreed that essential elements to communicate to patients before starting and periodically during opioid therapy include realistic expected benefits, common and serious harms, and expectations for clinician and patient responsibilities to mitigate risks of opioid therapy.

Clinicians should involve patients in decisions about whether to start or continue opioid therapy. Given potentially serious risks of long-term opioid therapy, clinicians should ensure that patients are aware of potential benefits of, harms of, and alternatives to opioids before starting or continuing opioid therapy. Clinicians are encouraged to have open and honest discussions with patients to inform mutual decisions about whether to start or continue opioid therapy. Important considerations include the following:

Be explicit and realistic about expected benefits of opioids, explaining that while opioids can reduce pain during short-term use, there is no good evidence that opioids improve pain or function with long-term use, and that complete relief of pain is unlikely (clinical evidence review, KQ1).

Emphasize improvement in function as a primary goal and that function can improve even when pain is still present.

Advise patients about serious adverse effects of opioids, including potentially fatal respiratory depression and development of a potentially serious lifelong opioid use disorder that can cause distress and inability to fulfill major role obligations.

Advise patients about common effects of opioids, such as constipation, dry mouth, nausea, vomiting, drowsiness, confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids. To prevent constipation

associated with opioid use, advise patients to increase hydration and fiber intake and to maintain or increase physical activity. Stool softeners or laxatives might be needed.

Discuss effects that opioids might have on ability to safely operate a vehicle, particularly when opioids are initiated, when dosages are increased, or when other central nervous system depressants, such as benzodiazepines or alcohol, are used concurrently.

Discuss increased risks for opioid use disorder, respiratory depression, and death at higher dosages, along with the importance of taking only the amount of opioids prescribed, i.e., not taking more opioids or taking them more often.

Review increased risks for respiratory depression when opioids are taken with benzodiazepines, other sedatives, alcohol, illicit drugs such as heroin, or other opioids.

Discuss risks to household members and other individuals if opioids are intentionally or unintentionally shared with others for whom they are not prescribed, including the possibility that others might experience overdose at the same or at lower dosage than prescribed for the patient, and that young children are susceptible to unintentional ingestion. Discuss storage of opioids in a secure, preferably locked location and options for safe disposal of unused opioids (188).

Discuss the importance of periodic reassessment to ensure that opioids are helping to meet patient goals and to allow opportunities for opioid discontinuation and consideration of additional nonpharmacologic or nonopioid pharmacologic treatment options if opioids are not effective or are harmful.

Discuss planned use of precautions to reduce risks, including use of prescription drug monitoring program information (see Recommendation 9) and urine drug testing (see Recommendation 10). Consider including discussion of naloxone use for overdose reversal (see Recommendation 8).

Consider whether cognitive limitations might interfere with management of opioid therapy (for older adults in particular) and, if so, determine whether a caregiver can responsibly co-manage medication therapy. Discuss the importance of reassessing safer medication use with both the patient and caregiver.

Given the possibility that benefits of opioid therapy might diminish or that risks might become more prominent over time, it is important that clinicians review expected benefits and risks of continued opioid therapy with patients periodically, at least every 3 months (see Recommendation 7).

Opioid Selection, Dosage, Duration, Follow-Up, and Discontinuation

4. When starting opioid therapy for chronic pain, clinicians should prescribe immediate-release opioids instead of extended-release/long-acting (ER/LA) opioids (recommendation category: A, evidence type: 4).

ER/LA opioids include methadone, transdermal fentanyl, and extended-release versions of opioids such as oxycodone, oxymorphone, hydrocodone, and morphine. The clinical evidence review found a fair-quality study showing a higher risk for overdose among patients initiating treatment with ER/LA opioids than among those initiating treatment with immediate-release opioids (77). The clinical evidence review did not find evidence that continuous, time-scheduled use of ER/LA opioids is more effective or safer than intermittent use of immediate-release opioids or that time-scheduled use of ER/LA opioids reduces risks for opioid misuse or addiction (KQ3).

In 2014, the FDA modified the labeling for ER/LA opioid pain medications, noting serious risks and recommending that ER/LA opioids be reserved for “management of pain severe enough to require daily, around-the-clock, long-term opioid treatment” when “alternative treatment options (e.g., nonopioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain” and not used as “as needed” pain relievers (121). FDA has also noted that some ER/LA opioids are only appropriate for opioid-tolerant patients, defined as patients who have received certain dosages of opioids (e.g., 60 mg daily of oral morphine, 30 mg daily of oral oxycodone, or equianalgesic dosages of other opioids) for at least 1 week (189). Time-scheduled opioid use can be associated with greater total average daily opioid dosage compared with intermittent, as-needed opioid use (contextual evidence review). In addition, experts indicated that there was not enough evidence to determine the safety of using immediate-release opioids for breakthrough pain when ER/LA opioids are used for chronic pain outside of active cancer pain, palliative care, or end-of-life care, and that this practice might be associated with dose escalation.

Abuse-deterrent technologies have been employed to prevent manipulation intended to defeat extended-release properties of ER/LA opioids and to prevent opioid use by unintended routes of administration, such as injection of oral opioids.

As indicated in FDA guidance for industry on evaluation and labeling of abuse-deterrent opioids (190), although abuse-deterrent technologies are expected to make manipulation of opioids more difficult or less rewarding, they do not prevent opioid abuse through oral intake, the most common route of opioid abuse, and can still be abused by nonoral routes. The “abuse-deterrent” label does not indicate that there is no risk for abuse. No studies were found in the clinical evidence review assessing the effectiveness of abuse-deterrent technologies as a risk mitigation strategy for deterring or preventing abuse. In addition, abuse-deterrent technologies do not prevent unintentional overdose through oral intake. Experts agreed that recommendations could not be offered at this time related to use of abuse-deterrent formulations.

In comparing different ER/LA formulations, the clinical evidence review found inconsistent results for overdose risk with methadone versus other ER/LA opioids used for chronic pain (KQ3). The contextual evidence review found that methadone has been associated with disproportionate numbers of overdose deaths relative to the frequency with which it is prescribed for chronic pain. In addition, methadone is associated with cardiac arrhythmias along with QT prolongation on the electrocardiogram, and it has complicated pharmacokinetics and pharmacodynamics, including a long and variable half-life and peak respiratory depressant effect occurring later and lasting longer than peak analgesic effect. Experts noted that the pharmacodynamics of methadone are subject to more inter-individual variability than other opioids. In regard to other ER/LA opioid formulations, experts noted that the absorption and pharmacodynamics of transdermal fentanyl are complex, with gradually increasing serum concentration during the first part of the 72-hour dosing interval, as well as variable absorption based on factors such as external heat. In addition, the dosing of transdermal fentanyl in mcg/hour, which is not typical for a drug used by outpatients, can be confusing. Experts thought that these complexities might increase the risk for fatal overdose when methadone or transdermal fentanyl is prescribed to a patient who has not used it previously or by clinicians who are not familiar with its effects.

Experts agreed that for patients not already receiving opioids, clinicians should not initiate opioid treatment with ER/LA opioids and should not prescribe ER/LA opioids for intermittent use. ER/LA opioids should be reserved for severe, continuous pain and should be considered only for patients who have received immediate-release opioids daily for at least 1 week. When changing to an ER/LA opioid for a patient previously receiving a different immediate-release opioid, clinicians should consult product labeling and reduce total daily dosage to account for incomplete opioid cross-tolerance. Clinicians should use additional caution with ER/LA opioids and consider a longer dosing interval when prescribing to patients with renal or hepatic dysfunction because decreased clearance of drugs among these patients can lead to accumulation of drugs to toxic levels and persistence in the body for longer durations. Although

there might be situations in which clinicians need to prescribe immediate-release and ER/LA opioids together (e.g., transitioning patients from ER/LA opioids to immediate-release opioids by temporarily using lower dosages of both), in general, avoiding the use of immediate-release opioids in combination with ER/LA opioids is preferable, given potentially increased risk and diminishing returns of such an approach for chronic pain.

When an ER/LA opioid is prescribed, using one with predictable pharmacokinetics and pharmacodynamics is preferred to minimize unintentional overdose risk. In particular, unusual characteristics of methadone and of transdermal fentanyl make safe prescribing of these medications for pain especially challenging.

Methadone should not be the first choice for an ER/LA opioid. Only clinicians who are familiar with methadone's unique risk profile and who are prepared to educate and closely monitor their patients, including risk assessment for QT prolongation and consideration of electrocardiographic monitoring, should consider prescribing methadone for pain. A clinical practice guideline that contains further guidance regarding methadone prescribing for pain has been published previously (191).

Because dosing effects of transdermal fentanyl are often misunderstood by both clinicians and patients, only clinicians who are familiar with the dosing and absorption properties of transdermal fentanyl and are prepared to educate their patients about its use should consider prescribing it.

5. When opioids are started, clinicians should prescribe the lowest effective dosage. Clinicians should use caution when prescribing opioids at any dosage, should carefully reassess evidence of individual benefits and risks when considering increasing dosage to ≥ 50 morphine milligram equivalents (MME)/day, and should avoid increasing dosage to ≥ 90 MME/day or carefully justify a decision to titrate dosage to ≥ 90 MME/day (recommendation category: A, evidence type: 3).

Benefits of high-dose opioids for chronic pain are not established. The clinical evidence review found only one study (84) addressing effectiveness of dose titration for outcomes related to pain control, function, and quality of life (KQ3). This randomized trial found no difference in pain or function between a more liberal opioid dose escalation strategy and maintenance of current dosage. (These groups were prescribed average dosages of 52 and 40 MME/day, respectively, at the end of the trial.) At the same time, risks for serious harms related to opioid therapy increase at higher opioid dosage. The clinical evidence review found that higher opioid dosages are associated with increased

210.0015555556

risks for motor vehicle injury, opioid use disorder, and overdose (KQ2). The clinical and contextual evidence reviews found that opioid overdose risk increases in a dose-response manner, that dosages of 50â€”

Reference

[Understanding Supervision and Assessment in Nursing](#)

[Statistical Theories of Mental Test Scores \(NA\)](#)