

The Mayo Clinic Diet: Eat well. Enjoy Life. Lose weight.

Susan Murphy Mayo Clinic researchers link ovarian cancer to bacteria colonization in microbiome

Share this:

ROCHESTER, Minn. — A specific colonization of microbes in the reproductive tract is commonly found in women with ovarian cancer, according to a new study from Mayo Clinic's Center for Individualized Medicine. The discovery, published in *Scientific Reports*, strengthens evidence that the bacterial component of the microbiome — a community of microorganisms that also consists of viruses, yeasts and fungi — is an important indicator for early detection, diagnosis and prognosis of ovarian cancer.

Abigail Asangba, Ph.D.

"In addition, we found a clear pattern that reveals women with early stage ovarian cancer have a significantly higher accumulation of the pathogenic microbes when compared to women with later-stage disease," says Abigail Asangba, Ph.D., a microbiome researcher within the Center for Individualized Medicine. "In later stages, the number of microbes fades. This strong signal could potentially help us diagnose women earlier and save lives — similar to how a noninvasive Pap smear is used to detect cervical cancer."

The study also suggests that a higher accumulation of pathogenic microbes plays a role in treatment outcomes and could be a potential indicator for predicting a patient's prognosis and response to therapy.

"We analyzed whether patients with similar outcomes also had a similar microbial composition before they started treatment, irrespective of stage, grade or histology of cancer, as well as other factors," Dr. Asangba says. "And we found that the patients with a higher accumulation of pathogenic microbes had poorer outcomes in comparison to those without."

Ovarian cancer ranks fifth in cancer deaths among women and is the second most common gynecological malignancy. An estimated 20,000 women in the U.S. are expected to be diagnosed with ovarian cancer in 2023, and nearly 13,000 will die from the disease, according to the American Cancer Society. Most women who are affected are usually diagnosed at an advanced stage because early-stage disease is usually asymptomatic. Furthermore, only 20% of cases are caused by genetic mutations, including BRCA1 and BRCA2 genes, while the remaining 80% of cases have no known cause.

Zeroing in on pathogenic microbes

For the study, the researchers investigated samples of 30 women undergoing a hysterectomy for ovarian cancer and compared them to samples of 34 women undergoing a hysterectomy for a benign condition. They used high-throughput sequencing to analyze the samples, which were recovered from the lower and upper reproductive tract, peritoneal fluid, urine, and anal microbiome.

P

In the women with ovarian cancer, the team observed a colonization of disease-causing bacteria, including Dialister, Corynebacterium, Prevotella and Peptoniphilus.

Marina R. Walther-Antonio, Ph.D.

"These microbes are known to be associated with other diseases, including other cancers, but more study is needed to know if they are a contributing driver of ovarian cancer," says Marina Walther-Antonio, Ph.D., a microbiome researcher within Mayo Clinic's Center for Individualized Medicine and a study author. Dr. Walther-Antonio is a member of the Mayo Clinic Comprehensive Cancer Center. She focuses on women's health, particularly gynecologic cancers.

"Our ultimate goal is to understand what role the microbiome plays in gynecologic cancers. We are exploring several potential avenues: the role in the causation of the disease, aggravation of the disease and treatment resistance," Dr. Walther-Antonio says.

The study is an extension of several other previously published studies by Dr. Walther-Antonio and her team that link the microbiome to endometrial cancer. In one study, the team found that a microbe called Porphyromonas somerae has a pathogenic role in endometrial cancer via intracellular activity.

Dr. Walther-Antonio says identifying microbiome signatures to predict the development of malignancies could lead to intervention before cancers have a chance to materialize.

"Our latest study provides a significant leap toward understanding the prognostic potential of the microbiome and places us a step closer to being able to help our patients," Dr. Walther-Antonio says.

Acknowledgements

This work was supported by a career development award from the Mayo Clinic Ovarian SPORC National Institutes of Health grant P50 CA136393), the Minnesota Ovarian Cancer Alliance and CTSA grant KL2TR002379 from the National Center for Advancing Translational Sciences.

###

About Mayo Clinic Comprehensive Cancer Center

Designated as a comprehensive cancer center by the National Cancer Institute, Mayo Clinic Comprehensive Cancer Center is defining new boundaries in possibility, focusing on patient-centered care, developing novel treatments, training future generations of cancer experts, and bringing cancer research to communities. At Mayo Clinic Comprehensive Cancer Center, a culture of innovation and collaboration is driving research breakthroughs that are changing approaches to cancer prevention, screening and treatment, and improving the lives of cancer survivors.

About Mayo Clinic

Mayo Clinic is a nonprofit organization committed to innovation in clinical practice, education and research, and providing compassion, expertise and answers to everyone who needs healing. Visit the Mayo Clinic News Network for additional Mayo Clinic news.

Media contact:

Susan Murphy, Mayo Clinic Communications, newsbureau@mayo.edu

Share this:

Related articles

Reference

[Crafting Phenomenological Research](#)

[Blue Marble Evaluation: Premises and Principles](#)